

The Blessing of Wealth The Curse of Poverty

Larry Blume & Klaus Ritzberger


Cornell University & IHS & The Santa Fe Institute



Introduction

- ▶ Why bats?
 - ▶ Gerald Wilkinson, “Reciprocal food sharing in the vampire bat.” *Nature* 308, 1984: 181–184.
 - ▶ —, “Food sharing in vampire bats.” *Scientific American* 262, 1990: 76-82.
- ▶ Sustaining cooperation?
 - ▶ David Kreps, Paul Milgrom, John Roberts and Robert Wilson, “Rational cooperation in the finitely repeated prisoners’ dilemma.” *JET* 27, 1982.
 - ▶ Robert Axelrod, *The Evolution of Cooperation*, 1984.
 - ▶ Elinor Ostrom, *Governing the Commons: The Evolution of Institutions for Collective Action*, 1990.

Model



- ▶ Two bats each eat 1 per day
- ▶ Each bat hunts once a day.
- ▶ A hunt returns 2 with pr p , or 0.
- ▶ Bats maintain an inventory.
- ▶ A bat **may** share 1 with the other bat.
- ▶ A bat dies when she fails to eat.

The inventory space is \mathcal{I} , which describes each bat's current inventory.

$b_i = -1$ is **death**, an absorbing state.



Model

- ▶ Let Ω denote the sample space on which the processes are built: $\omega_t = (\omega_{1t}, \omega_{2t})$ where each ω_{it} describes the outcome of i 's date- t hunt, **s**uccess or **f**ailure.
- ▶ From each strategy profile (σ_1, σ_2) and the initial inventory $b = (b_{10}, b_{20})$ compute $\tau_b^i(\omega) = \inf\{t : b_{it} = -1\}$, the time at which i dies.
- ▶ Bat i 's payoff function is

$$u_i(\sigma_1, \sigma_2) = \sum_{t=0}^{\tau_b^i(\omega)} \delta^t.$$



Dynamics

- ▶ An **action** for bat i is a choice to **share** S or **withhold** W 1 unit from bat j . $\mathcal{A} = \{S, W\}$.
- ▶ A **state** $q \in Q$ of the game is a quadruple $q = (b_1, b_2, \omega_1, \omega_2)$ where $(b_1, b_2) \in \mathcal{I}$ are the bat's inventory levels, and ω_i describes the outcome of bat i 's hunt. State q_t describes the date- t physical situation after hunting.

Taking $s = S = 1$ and $f = W = 0$, the dynamics are


$$\begin{array}{ccc} \longrightarrow & \begin{array}{c} b_{1t} \\ b_{2t} \\ \underbrace{\hspace{1.5cm}} \\ \mathbf{b}_t \end{array} & \rightsquigarrow \begin{array}{c} b_{1t}, \omega_{1t} \\ b_{2t}, \omega_{2t} \\ \underbrace{\hspace{1.5cm}} \\ \mathbf{q}_t \end{array} & \longrightarrow & \begin{array}{c} b_{1t} + \omega_{1t} - a_{1t} + a_{2t} - 1 \\ b_{2t} + \omega_{2t} - a_{2t} + a_{1t} - 1 \\ \underbrace{\hspace{3.5cm}} \\ \mathbf{b}_{t+1} \end{array} & \rightsquigarrow \end{array}$$

Strategies


- ▶ A **rule** is a map $r : Q \rightarrow \mathcal{A}$. \mathcal{R} denotes the set of rules.
- ▶ A **partial history** is a structure $(q_0, a_{10}, a_{20}, \dots, q_t, a_{1t}, a_{2t})$.
Let \mathcal{H} denote the set of **partial histories**. $h_t \in \mathcal{H}$ is the sequence of states and actions **through** date t .
- ▶ A **strategy** for bat i is a map $\sigma : \mathcal{H} \rightarrow \mathcal{R}$ where $\sigma(h_{t-1})$ is the rule employed at date t .

Some Rules


- ▶ **Autarchy.** Each bat contributes to and withdraws from her own inventory. No sharing takes place.



- ▶ **Simple Sharing.** A successful bat shares with an unsuccessful bat.



- ▶ **Wealth-Based Sharing.** A successful bat shares with the other bat if and only if she is wealthier than the other bat.



We only allow rules that **share** (or not) **upon success**.

Questions

The possibilities for cooperation:

- ▶ Is sharing over some or all of the state space optimal?

Yes, on all of \mathcal{I} .

- ▶ For large δ , are there equilibria which support sharing on some or all of the state space?

Depends on p , and only for some part of \mathcal{I} .

- ▶ For large δ , are there equilibria which achieve the welfare optima?

No.

Folk Theorems

These questions are normally answered by folk theorems.

Requirements for the folk theorem:

- ▶ The set of feasible long-run average payoffs is state-independent.
- ▶ The long-run average min-max payoffs are state-independent.
- ▶ The dimension of the set of long-run average feasible payoffs is 2.

In our game.

- ▶ For $p < 1/2$ the only feasible long-run average payoff is 0.
- ▶ For $p > 1/2$, the maximal long-run average payoff is state-dependent.

Value Functions

- ▶ Strategies determine a stochastic process on \mathcal{I} .
- ▶ τ_b^i is the first time the process hits -1 .
- ▶ The value of being at $b \in \mathcal{I}$ is

$$V_i(b) = \mathbb{E} \left\{ \sum_{t=0}^{\tau_b^i(\omega)} \delta^t \right\} = \frac{1 - \delta \mathbb{E} \left\{ \delta^{\tau_b^i(\omega)} \right\}}{1 - \delta}$$

$$\lim_{\delta \uparrow 1} V_i(b) = \begin{cases} 1 + \mathbb{E} \left\{ \tau_b^i(\omega) \right\} & \text{if } \text{pr} \left\{ \tau_b^i(\omega) < \infty \right\} = 1, \\ \infty & \text{otherwise} \end{cases}$$

- ▶ The average discounted value at $b \in \mathcal{I}$ is

$$ADV_i(b) = 1 - \delta \mathbb{E} \left\{ \delta^{\tau_b^i(\omega)} \right\},$$

$$\lim_{\delta \uparrow 1} ADV_i(b) = \text{pr} \left\{ \tau_b^i(\omega) = \infty \right\}.$$

The value to bat i of being in state (b_1, b_2) , by recursion:

$$V_1^{aut}(b_1) = \begin{cases} 1 + \delta p V_1^{aut}(b_1 + 1) + \delta(1 - p) V_1^{aut}(b_1 - 1) & \text{for } b \geq 1, \\ p + \delta p V_1^{aut}(b_1 + 1) & \text{for } b = 0. \end{cases}$$

This is a **linear second-order difference equation** with two boundary conditions: $V_1^{aut}(-1) = 0$ and $\lim_{b_1 \rightarrow \infty} V_1^{aut}(b_1) = 1/(1 - \delta)$.

$$V_1^{aut}(b_1) = \frac{1}{1 - \delta} \left(1 - \frac{\mu^{b_1+1}}{\delta} \right)$$

Proofs

So how does one compute these stopping times?

For, at $b_1 > b_2 > 0$ substituting (19) into the defining equation yields by (10)

$$\begin{aligned}w(b_1, b_2) &= 1 - \delta + \delta p^2 [1 - \lambda_2^{b_2+1} + \lambda_2^{b_2+1} w(b_1 - b_2, 0)] \\&\quad + 2\delta p(1-p) [1 - \lambda_2^{b_2} + \lambda_2^{b_2} w(b_1 - b_2, 0)] \\&\quad + \delta(1-p)^2 [1 - \lambda_2^{b_2-1} + \lambda_2^{b_2-1} w(b_1 - b_2, 0)] \\&= 1 - (\delta p^2 \lambda_2^2 + 2\delta p(1-p)\lambda_2 + \delta(1-p)^2) \lambda_2^{b_2-1} [1 - w(b_1 - b_2, 0)] \\&= 1 - \lambda_2^{b_2} [1 - w(b_1 - b_2, 0)] = 1 - \lambda_2^{b_2} + \lambda_2^{b_2} w(b_1 - b_2, 0)\end{aligned}$$

and at $b_2 \geq b_1 > 0$ substituting (20) into the defining equation and using (10) yields

$$\begin{aligned}w(b_1, b_2) &= 1 - \delta + \delta p^2 [1 - \lambda_2^{b_1+1} + \lambda_2^{b_1+1} w(0, b_2 - b_1)] \\&\quad + 2\delta p(1-p) [1 - \lambda_2^{b_1} + \lambda_2^{b_1} w(0, b_2 - b_1)] \\&\quad + \delta(1-p)^2 [1 - \lambda_2^{b_1-1} + \lambda_2^{b_1-1} w(0, b_2 - b_1)] \\&= 1 - (\delta p^2 \lambda_2^2 + 2\delta p(1-p)\lambda_2 + \delta(1-p)^2) \lambda_2^{b_1-1} [1 - w(0, b_2 - b_1)] \\&= 1 - \lambda_2^{b_1} + \lambda_2^{b_1} w(0, b_2 - b_1),\end{aligned}$$

as required. Substituting (19) and (1) into the defining equation for $w(b_1, 0)$ and using (3), (16), and (18) yields for even $b_1 > 1$

$$\begin{aligned}w(b_1, 0) &= 1 - \delta + \delta p^2 [1 - \lambda_2 + \lambda_2 w(b_1, 0)] + \delta p(1-p) [1 - \lambda_2 + \lambda_2 w(b_1 - 2, 0)] \\&\quad + \delta(1-p)^2 \left(1 - \frac{\mu_2^{b_1}}{\delta}\right) + \delta p(1-p) \left(1 - \frac{\mu_2^{b_1+2}}{\delta}\right) \\&= 1 - \delta p \lambda_2 + \delta p^2 \lambda_2 w(b_1, 0) + \delta p(1-p) \lambda_2 w(b_1 - 2, 0) - (1-p) \frac{\mu_2^{b_1+1}}{\delta} \\&= 1 - \delta p \lambda_2 - (1-p) \frac{\mu_2^{b_1+1}}{\delta} \\&\quad + \delta p^2 \lambda_2 \left[1 - B p \mu_2^{b_1+1} - (1 - B \mu_2 - w(0, 0)) \gamma^{\frac{b_1}{2}}\right] \\&\quad + \delta p(1-p) \lambda_2 \left[1 - B \mu_2^{b_1-1} - (1 - B \mu_2 - w(0, 0)) \gamma^{(b_1-2)/2}\right] \\&= 1 - \left[(1-p) \frac{\mu_2^2}{\delta} + \delta p^2 \lambda_2 B \mu_2^2 + \delta p(1-p) \lambda_2 B\right] \mu_2^{b_1-1} \\&\quad - \delta p^2 \lambda_2 [1 - B p \mu_2 - w(0, 0)] \gamma^{b_1/2} \\&\quad - \delta p(1-p) \lambda_2 [1 - B \mu_2 - w(0, 0)] \gamma^{(b_1-2)/2} \\&= 1 - \left[\frac{1-p}{\delta} + \frac{p(1-p)\lambda_2}{\delta(\mu_2 - p\lambda_2)}\right] \mu_2^{b_1-1} - [1 - B \mu_2 - w(0, 0)] \gamma^{b_1/2} \\&= 1 - \frac{(1-p)\mu_2}{\delta(\mu_2 - p\lambda_2)} \mu_2^{b_1+1} - [1 - B \mu_2 - w(0, 0)] \gamma^{b_1/2} \\&= 1 - B \mu_2^{b_1+1} - [1 - B \mu_2 - w(0, 0)] \gamma^{b_1/2}\end{aligned}$$

Welfare

Take as a welfare function $W(b_1, b_2)$ the sum of the bats' expected lifetimes.

- ▶ It can be described as the fixed point of a Bellman operator.
- ▶ W is symmetric around the diagonal.
- ▶ W is strictly increasing.
- ▶ If $b_1 > b_2$, then $W(b_1, b_2) < W(b_1 + 1, b_2 - 1)$. Moving diagonally towards the main diagonal is welfare improving.

Results

- ▶ The welfare-optimal strategy is **wealth sharing** whenever both bats are alive.

We look for equilibria with a **grim trigger**. A successful bat may defect from sharing and revert to autarchy but with one more unit.

- ▶ Autarky everywhere is an equilibrium **for any** p .
- ▶ Autarky is the **only** equilibrium for $p < 1/2$.

Proof. $V_1^{aut}(b_1 + 1) + V_2^{aut}(b_2 + 1) > W(b_1, b_2)$.

But if there is an equilibrium other than autarchy, then for all i ,

$$V_i^{eq}(b_1, b_2) \geq V_i^{aut}(b_i + 1).$$

Results

- ▶ **Bilateral sharing** on all of \mathcal{I} is **not** an equilibrium.

Proof. Diagonals are invariant. The incentive constraint is violated at $\min\{b_1, b_2\} = 0$.

- ▶ If $p > 1/2$ and δ is sufficiently near 1, then **bilateral sharing** on the interior of \mathcal{I} **is** an equilibrium.
- ▶ If $p > 1/2$ and δ sufficiently near 1, there is an equilibrium in which the **wealth-sharing** rule **is** used on the set $\{(b_1, b_2) : |b_1 - b_2| \leq 1, b_1, b_2 > 0\}$.

Summary

- ▶ Sharing is not possible in poor societies, $p < 1/2$.
- ▶ Sharing is possible in wealthy societies, $p > 1/2$, and more so for wealthier societies.
- ▶ Nonetheless, even in wealthy societies the welfare optimum cannot be achieved.

